

RESEARCH ARTICLE

Open Access



The *Candida* species that are important for the development of atrophic glossitis in xerostomia patients

Sachika Nakamura^{1,2}, Mariko R. Okamoto^{1,2,3}, Ken Yamamoto^{1,4}, Akihisa Tsurumoto⁴, Yoko Yoshino^{1,5}, Hiroshi Iwabuchi⁶, Ichiro Saito^{1,7}, Nobuko Maeda² and Yoichi Nakagawa^{1*}

Abstract

Background: The purpose of this study was to clarify the species of *Candida* that are important for the development of atrophic glossitis in xerostomia patients.

Methods: A total of 231 patients with subjective dry mouth were enrolled in the present study. Logistic regression analysis was performed to clarify the contribution of each *Candida* species and other variables to the development of atrophic glossitis. The dependent variable was the absence/presence of atrophic glossitis. The *Candida* colony-forming units (CFU) of *C. albicans*, *C. glabrata*, *C. tropicalis*, and *C. krusei*, as well as age, gender, resting (RSFR) and stimulated (SSFR) whole salivary flow rate, and denture-wearing status, were treated as explanatory variables.

Results: Logistic regression analysis showed that two factors were closely associated with the presence of atrophic glossitis: an increase in *C. albicans* CFU and a decrease in the SSFR.

Conclusions: *C. albicans*, but not non-*albicans Candida*, was associated with atrophic glossitis in xerostomia patients who had no systemic predisposing factors, indicating that *C. albicans* remains a treatment target for *Candida*-related atrophic glossitis.

Keywords: Atrophic glossitis, *Candida* species, Candidiasis, Xerostomia, Logistic regression analysis

Background

Oral candidiasis is sub-classified into three major variants: pseudomembranous, erythematous, and hypertrophic [1, 2]. In addition to these variants, angular cheilitis, median rhomboid glossitis, and denture stomatitis have been recognized as *Candida*-associated lesions; moreover, *Candida* super-infection in oral lichen planus is frequently encountered. The various forms of oral candidiasis are caused by interactions between the host defenses and fungal virulence factors.

Atrophic glossitis is caused by complete or partial lingual papillary atrophy. It exhibits a smooth, glossy appearance with a red or pink background and is primarily a manifestation of an underlying condition. Nutritional

deficiencies (iron, folic acid, vitamin B₁₂, riboflavin, and niacin) are the most common causes [3]. Other etiological factors of atrophic glossitis include hyposalivation and candidiasis [2, 4–6]. An increase in *Candida* colony counts, a low salivary flow rate, and advanced age were identified as being risk factors that were closely associated with the development of atrophic glossitis [7, 8]. However, the *Candida* species that are the most relevant to the development of atrophic glossitis remain unclear.

Candida albicans is the primary causative organism of oral candidiasis. Recent studies have suggested that non-*albicans Candida* (NAC) species, such as *C. glabrata*, *C. parapsilosis*, *C. tropicalis*, and *C. krusei* are also pathogenic in humans [2, 8–13]. One of these studies showed that *C. albicans* was frequently isolated from the tongue, while *C. glabrata* was most frequently isolated from the angle of the mouth, suggesting that certain *Candida* species are associated with oral candidiasis at particular sites [8]. In a study on the relationship between *Candida*

* Correspondence: nakagawa-y@tsurumi-u.ac.jp

¹Department of Clinical Pathophysiology, Tsurumi University School of Dental Medicine, Tsurumi University Dental Hospital, 2-1-3 Tsurumi, Tsurumi-ku, Yokohama 230-8501, Japan

Full list of author information is available at the end of the article



species and local and systemic predisposing factors, only *C. albicans* infections were significantly associated with the use of inhaled steroids and antibiotics and superinfection with oral lichen planus, whereas the presence of removable dentures was significantly associated with the detection of NAC or a mixed flora consisting of *C. albicans* and NAC [9]. Similarly, the incidence of colonization with more than one *Candida* species was found to be higher in cases of denture stomatitis than in cases involving other forms of oral candidiasis [10]. *C. albicans* and *C. glabrata* were the most prevalent species detected in cases of mixed infections [10, 14]. However, no association has been confirmed between a specific *Candida* species and any particular sign or symptom of oral candidiasis.

The purpose of this study was to clarify the species of *Candida* that are important for the development of atrophic glossitis in xerostomia patients.

Methods

Patient selection

The subjects enrolled in the study were patients at the Dry Mouth Clinic at Tsurumi University Dental Hospital. Patients who had subjective dry mouth (xerostomia) were examined to determine their salivary flow rate and oral findings, and a mycological examination was performed. The examinations were performed between January 2010 and February 2013. Patients with systemic predisposing factors for candidiasis, such as diabetes mellitus, chronic obstructive pulmonary disease, immunodeficiency, and malnutrition, and patients who were using medications that predisposed them to candidiasis, such as antibiotics and steroids, were excluded from this study. A total of 231 patients (mean age \pm standard deviation, 66.6 ± 12.6 years; range 33 to 96 years), were enrolled in this study. Forty-two of the patients were men (18.2%), and 189 were women (81.8%).

Evaluation of atrophic glossitis

The presence of atrophic glossitis was determined by the attending dentist. The severity and extent of papillary atrophy and loss were graded from 0 to 4 as follows: 0, none; 1, mild; 2, moderate; 3, severe; and 4, profound loss (Fig. 1). Atrophic glossitis was considered mild when the lingual papillae in the affected area of the tongue were smaller in size than in the intact area. Partial loss (<50%) of the lingual papillae of the dorsum of the tongue was considered to indicate moderate disease, and partial loss (>50%) was considered to indicate severe disease. Patients who showed loss of the lingual papillae throughout the dorsum of the tongue were considered to have profound loss. When the attending dentist could not determine the amount of papillary atrophy, a consensus was reached by discussion with an oral surgeon (Y.N.) with 30 years of experience [7].




Fig. 1 The grades of atrophic glossitis. 0, none; 1 (mild), the lingual papillae in the affected area of the tongue were smaller in size than those in the intact area; 2 (moderate), partial loss (<50%) of the lingual papillae on the dorsum of the tongue; 3 (severe), partial loss (>50%) of the lingual papillae on the dorsum of the tongue; and 4 (profound loss)

Salivary secretion test

To measure the resting whole salivary flow rate (RSFR), each subject sat at rest, and saliva was collected into a cup during a 15-min period in the absence of masticatory movements. The saliva volume was measured using a disposable syringe. An RSFR of ≤ 1.5 mL/15 min (≤ 0.1 mL/min) was classified as hyposalivation according to previously determined criteria [15, 16]. The stimulated whole salivary flow rate (SSFR) was also measured by collecting saliva during a 10-min period. Salivation was stimulated by chewing a piece of gum (Free zone gum; Lotte Co., Ltd., Tokyo, Japan). An SSFR of ≤ 10 mL/10 min was classified as hyposalivation.

Fungal culture

The dorsum of the tongue was swabbed 10 times with a cotton applicator (Eiken Chemical Co., Ltd., Tokyo, Japan), and the swab was directly inoculated onto CHROMagar™ *Candida* selective plates (CHROMagar Microbiology, Paris, France) [7, 17]. The number of *Candida* colonies was counted after incubation at 30 °C for 48 h and was expressed as colony-forming units (CFU) per plate. Each *Candida* species was identified by colony color in accordance with the manufacturer's manual [17].

Statistical analyses

A multivariate logistic regression analysis was used to clarify the contribution of each *Candida* species and the other variables to the development of atrophic glossitis. The

dependent variable was the absence (Grade 0)/presence (Grades 1–4) of atrophic glossitis, while the *Candida* CFU of *C. albicans*, *C. glabrata*, *C. tropicalis*, and *C. krusei* as well as age, RSFR, SSFR and denture-wearing status were treated as explanatory variables. The results of the logistic regression analysis were used to ascertain which independent variables had a major effect on the absence or presence of atrophic glossitis.

Salivation and the patient's denture-wearing status were analyzed as factors associated with increased *Candida* colony counts. To determine the effects of hyposalivation and denture wearing on *Candida* colony counts, differences among the groups were evaluated by the Kruskal-Wallis H test, followed by multiple comparison testing using the Mann-Whitney U test with Bonferroni correction. A total of four groups were formed through combinations of two factors: salivation status (hyposalivation [SSFR ≤ 10 mL/10 min] or normosalivation [SSFR > 10 mL/10 min]) and denture-wearing status (denture wearing/no dentures).

All statistical analyses were performed using the SPSS software program, ver. 22.0 J (IBM Japan, Tokyo). *P* values of < 0.05 were considered to indicate statistical significance.

Results

Study population

Overall, atrophic glossitis was noted in 46 patients (Table 1), tongue redness in 58, palate redness in 16, buccal mucosa redness in 10, lip redness in 16 and angular cheilitis in 25. The mean RSFR was 2.1 ± 2.7 mL/15 min (range, 0–16.0 mL/15 min); 26 (11.3%) patients had an RSFR of 0 mL/15 min, and 141 (61.0%) patients had an RSFR of ≤ 1.5 mL/15 min. The mean SSFR was 11.3 ± 7.0 mL/10 min (range, 0.6–40.6 mL/10 min); 109 (47.2%) patients had an SSFR ≤ 10 mL/10 min. According to the Japanese criteria [18], 26 (11.3%) patients were diagnosed with Sjögren's syndrome. Ninety-two (39.8%) patients wore dentures. Partial removable dentures were more commonly used than full dentures, but we did not consider the type of denture in our analysis.

Table 1 Characteristics of study population

Atrophic glossitis	Number	Age	Male:Female	Saliva flow rate		Denture wearer
				RSFR	SSFR	
Grade 0	185	63.4 ± 12.5	37:148	2.4 ± 2.7	12.2 ± 6.9	67/185 (36.2%)
Grade 1	15	72.8 ± 10.8	1:14	1.9 ± 3.3	11.3 ± 6.1	8/15 (53.3%)
Grade 2	23	68.2 ± 11.9	2:21	1.0 ± 2.4	6.3 ± 6.2	11/23 (47.8%)
Grade 3	5	74.4 ± 10.0	2:3	1.1 ± 0.7	7.7 ± 4.8	4/5 (80.0%)
Grade 4	3	83.0 ± 6.5	0:3	0.1 ± 0.1	3.9 ± 0.8	2/3 (66.7%)
Total	231	66.6 ± 12.6	42:189	2.1 ± 2.7	11.3 ± 7.0	92/231(39.8%)

RSFR resting whole salivary flow rate (mL/15 min), SSFR stimulated whole salivary flow rate (mL/10 min); Values are mean \pm SD

Candida colonization according to the severity of atrophic glossitis

One hundred thirty-five (58.4%) patients were *Candida*-positive (Table 2). The species of *Candida* in these cases were as follows: *C. albicans*, $n = 119$ (88.1%); *C. glabrata*, $n = 50$ (37.0%); *C. tropicalis*, $n = 20$ (14.8%), and *C. krusei*, $n = 9$ (6.7%). *C. albicans* and *C. glabrata* were simultaneously detected in 41 (30.4%) of the 135 patients.

The variables of patients with and without atrophic glossitis

When the participants were divided into two groups according to the absence/presence of atrophic glossitis (Table 3), the number of *C. albicans* CFU in the group with atrophic glossitis was found to be significantly higher (255.6 ± 750.0 CFU) than that in the group without atrophic glossitis (35.2 ± 92.3 CFU) ($P < 0.001$). The *C. glabrata* CFU in the group with atrophic glossitis (37.9 ± 115.2 CFU) was significantly lower than that in the group without atrophic glossitis (45.8 ± 278.7 CFU) ($P < 0.001$).

In the group with atrophic glossitis, the mean RSFR was 1.3 ± 2.6 mL/15 min, while the mean SSFR was 7.9 ± 6.4 mL/10 min. In contrast, in the group without atrophic glossitis, the mean RSFR was 2.4 ± 2.7 mL/15 min, and the mean SSFR was 12.2 ± 6.9 mL/10 min. Thus, both the mean resting ($P < 0.001$) and stimulated ($P < 0.001$) saliva flow rates were significantly lower in the patients with atrophic glossitis.

A chi-squared test revealed that the gender of the groups with and without atrophic glossitis did not differ to a statistically significant extent ($P = 0.211$; Table 4). The chi-squared test also revealed that denture wearing was associated with atrophic glossitis ($P = 0.025$).

Correlations among the colonizing species of *Candida*, age, and salivary flow rate

Significant relationships were observed between age and the amounts of *Candida* species as well as between age and the RSFR according to the Spearman's rank correlation coefficients (Table 5). Significant relationships were also observed among the *Candida* species, except between

Table 2 Prevalence of *Candida* colonization according to the severity of atrophic glossitis

Atrophic glossitis	Detection of <i>Candida</i> ^a				
	<i>C. albicans</i>	<i>C. glabrata</i>	<i>C. tropicalis</i>	<i>C. krusei</i>	<i>Candida</i> spp.
Grade 0	76/185	30/185	16/185	5/185	14/185
Grade 1	13/15	4/15	3/15	0/15	0/15
Grade 2	22/23	11/23	1/23	4/23	3/23
Grade 3	5/5	2/5	0/5	0/5	0/5
Grade 4	3/3	3/3	0/3	0/3	1/3
Total	119/231	^b 50/231	20/231	9/231	18/231

^aThe culture test was positive in 135/231 (58.4%) patients. Values are number of cases

^b*C. albicans* and *C. glabrata* were simultaneously detected in 41 cases

C. albicans and *C. krusei*, which showed weak relationships. Negative correlations were observed between the amounts of *Candida* species and the salivary flow rate. There was no significant correlation between *C. krusei* and the salivary flow rate.

Factors associated with atrophic glossitis

Logistic regression analysis determined the factors that were closely associated with the presence of atrophic glossitis (Table 6). These factors included an increase in *C. albicans* CFU (Odds ratio, 1.004; 95% CI, 1.001–1.007) and a decrease in SSFR (Odds ratio, 0.919; 95% CI, 0.845–0.999); *C. albicans* showed the greatest contribution to atrophic tongue. Although *C. glabrata* was detected in 50 of 135 (37.0%) patients and was the second-most common species after *C. albicans* (Table 2), *C. glabrata* was not significantly associated with the presence of atrophic glossitis (Table 6).

The influences of the stimulated salivary flow rate and denture wearing on the amounts of *Candida* colonies

Logistic regression analysis demonstrated that denture wearing was not associated with atrophic glossitis (Table 6).

Table 3 Comparison of variables between absent (Grade 0) and present (Grade1~4) groups of atrophic glossitis

	Atrophic glossitis		P-value
	Grade 0 n = 185	Grade 1~4 n = 46	
<i>C. albicans</i> (CFU)	35.2 ± 92.3	255.6 ± 750.0	<0.001
<i>C. glabrata</i> (CFU)	45.8 ± 278.7	37.9 ± 115.2	<0.001
<i>C. tropicalis</i> (CFU)	0.6 ± 5.8	3.9 ± 18.6	0.058
<i>C. krusei</i> (CFU)	0.9 ± 6.3	0.3 ± 1.2	0.974
RSFR (mL/15 min)	2.4 ± 2.7	1.3 ± 2.6	<0.001
SSFR (mL/10 min)	12.2 ± 6.9	7.9 ± 6.4	<0.001
Age	65.4 ± 12.5	71.4 ± 12.0	0.004

Values are expressed as Mean ± SD. P-value, Mann-Whitney U-test

RSFR resting whole salivary flow rate, SSFR stimulated whole salivary flow rate

Table 4 Comparison of gender and denture-wearing status between absent (Grade 0) and present (Grade1~4) groups of atrophic glossitis

		Atrophic glossitis		Total	P-value
		Grade 0	Grade 1~4		
Gender	Male	37	5	42	0.221
	Female	148	41	189	
	Total	185	46	231	
Denture	Wearing	67	25	92	0.025
	No dentures	118	21	139	
	Total	185	46	231	

Values are number of cases. P-value, Chi square test

RSFR resting whole salivary flow rate, SSFR stimulated whole salivary flow rate

The question then arose as to whether denture wearing promoted an increase in the number of *C. albicans* CFU. We therefore examined the relationships between hyposalivation and denture wearing and the amount of *C. albicans* using the Kruskal-Wallis H test, followed by multiple comparison testing using the Mann-Whitney U test with Bonferroni correction (Fig. 2). Both denture-wearing (mean CFU ± standard deviation, 199.6 ± 737.0 CFU) and non-denture-wearing hyposalivation patients (82.5 ± 173.1 CFU) showed a significantly high number of *C. albicans* colonies compared with the non-denture-wearing normosalivation patients (13.8 ± 42.4 CFU). There was no significant difference in the number of *C. albicans* colonies between the denture-wearing (64.6 ± 167.6 CFU) and non-denture-wearing normosalivation patients. Similarly, no significant difference in *C. albicans* colony counts was noted between the denture-wearing and non-denture-wearing hyposalivation patients.

Discussion

The purpose of this study was to clarify the species of *Candida* that are most strongly associated with the development of *Candida*-associated atrophic glossitis because recent studies have suggested the importance of NAC pathogenesis in oral candidiasis. Our logistic regression analysis showed an association between increased amounts of *C. albicans* colonies and atrophic glossitis. However, no significant relationship between NAC and atrophic glossitis was observed.

With regard to the methods that are used to obtain samples from the oral cavity, although an oral rinse method is helpful for quantification, this method does not evaluate localized sites of infection [19]. In the present study, swab testing was used to evaluate local colonization on the dorsum of the tongue. Swab testing is simple to perform and is thought to be useful for quantitative estimation since the results obtained from rinsing and swab methods have been found to be correlated [20, 21]. A chromogenic media-

Table 5 Correlations among amount of *Candida* species, age, and salivary flow rate

	<i>C. albicans</i>	<i>C. glabrata</i>	<i>C. tropicalis</i>	<i>C. krusei</i>	RSFR	SSFR
Age	0.296**	0.288**	0.175**	0.155*	-0.210**	-0.099
<i>C. albicans</i>		0.345**	0.157*	0.039	-0.430**	-0.351**
<i>C. glabrata</i>			0.189**	0.246**	-0.320**	-0.269**
<i>C. tropicalis</i>				0.178**	-0.248**	-0.278**
<i>C. krusei</i>					-0.079	-0.031
RSFR						0.701**
SSFR						

Values represent Spearman's rank correlation coefficients; * $P < 0.05$, ** $P < 0.001$
 RSFR resting whole salivary flow rate, SSFR stimulated whole salivary flow rate


based commercial system, CHROMagar™ *Candida*, was used for quantitative estimation; this selective and differential medium is used for the isolation and identification of *Candida* species [19]. The particular advantage of this system is its ability to detect mixed yeast infections in clinical samples [22]. Studies have suggested that *C. albicans*, *C. tropicalis*, *C. glabrata*, and *C. krusei* can be identified accurately using CHROMagar™ *Candida* [23–26]. Although it has been reported that the dark green appearance of *C. dubliniensis* can also be reliably distinguished from that of *C. albicans* [27], a more detailed method, such as molecular fingerprinting, is needed to discriminate between these two species since *C. albicans* and *C. dubliniensis* share many phenotypic characteristics [28]. In the present study, no molecular methods were applied; thus, *C. dubliniensis* colonies could not be distinguished from *C. albicans* on CHROMagar™ *Candida* plates. However, *C. dubliniensis* is an opportunistic oral pathogen that is typically isolated from patients infected with human immunodeficiency virus (HIV) [29]. Since the present study did not include any such patients, there is little concern about the potential for confusion.

Table 6 Predictors of atrophic glossitis -Multivariate logistic regression analysis-

	Coefficient	P-value	Odds ratio	95% CI	
				Lower	Upper
Age	0.029	0.135	1.029	0.991	1.069
Gender	0.804	0.196	2.234	0.661	7.549
<i>C. albicans</i> (CFU)	0.004	0.004	1.004	1.001	1.007
<i>C. glabrata</i> (CFU)	-0.001	0.366	0.999	0.996	1.001
<i>C. tropicalis</i> (CFU)	0.015	0.353	1.015	0.983	1.048
<i>C. krusei</i> (CFU)	-0.055	0.581	0.946	0.778	1.151
RSFR (mL/15 min)	0.045	0.675	1.047	0.846	1.295
SSFR (mL/10 min)	-0.085	0.048	0.919	0.845	0.999
Denture	0.364	0.389	1.439	0.629	3.294
Constant	-3.752	0.010	0.023		

RSFR resting whole salivary flow rate, SSFR stimulated whole salivary flow rate
 Gender; female: 1, male: 0, Denture; denture wearing: 1, no dentures: 0

C. albicans, but not NAC, proved to be associated with atrophic glossitis in the present study. This result was consistent with the observations reported by Terai et al. [5, 30], who found that *C. albicans* is predominantly detected as a single infection. The detection of NAC or a mixed flora consisting of *C. albicans* and NAC is therefore limited in patients with atrophic glossitis. Although *C. glabrata* was detected in 50 of the 135 (37.0%) patients with *Candida*, the second-most common isolated species after *C. albicans* (Table 2), no significant association was observed between *C. glabrata* and atrophic glossitis (Table 6). Since the clinical manifestation of erythematous candidiasis is related to the proteinase production capacity of *C. albicans* [31], a low level of protease secretion might explain why *C. glabrata* did not contribute to atrophic glossitis.



The virulence factors of *Candida* include adherence, evasion of the host defenses, and the invasion and destruction of host tissue [1]. Hyphae penetrate perpendicularly, traversing the surface epithelium up to the spinous cell layer [32]. The tip of the penetrating hyphae is known to be rich in proteinases, lipases, and several other enzymes. *C. glabrata* is unable to grow in filamentous forms and cannot invade the epithelial layer [33]. In contrast to other *Candida* species, the amounts of secreted aspartyl proteinases (SAPs) and phospholipases (PLs) produced by *C. glabrata* are extremely low [11, 34]. In *C. albicans* and *C. glabrata* mixed infections, *C. albicans* promotes the penetration of *C. glabrata* into the oral epithelium as *C. glabrata* cannot do so alone [33]. Once *C. albicans* penetrates the epithelial layer, the final step in the infection process is damage, which is characterized by the loss of the superficial epithelium [35]. *C. albicans* induces both apoptosis and necrosis in the oral epithelial cells [36]. In a study using reconstituted human oral epithelia, the hyphal elements of *C. albicans* invaded and led to marked disorganization of the epithelium [33]. In contrast, *C. glabrata* induced less epithelial damage [33]. Different levels of epithelial loss, namely, the severity of atrophic glossitis, may be related to the levels of disorganization of the epithelium. The removal of the keratin layer by oral dynamics, i.e., the movement of the tongue and other muscles, might be enhanced in certain situations, especially in patients with hyposalivation, due to the reduced lubricant function of the saliva. Erythematous candidiasis is the most prevalent variant in patients with hyposalivation [37]. Furthermore, in the present study, a logistic regression analysis revealed that a low SSFR was associated with atrophic glossitis (Table 6).

High vascularity is a characteristic finding of erythematous candidiasis in addition to epithelial atrophy or a lack of a keratinized epithelial surface layer [32]. One hypothesis regarding the mechanism underlying these findings is that erythematous candidiasis is accompanied by the activation of a partially reactive defense mechanism and that it may represent a clinical expression in response to candidal antigens [38, 39]. Because patients with systemic immunosuppressive diseases were excluded from the present study, the oral mucosa in the patients was presumed to be immunocompetent. Atrophic glossitis may be induced even in patients with normal immunity; as such, factors other than immunosuppression, such as hyposalivation, are necessary for the establishment of atrophic glossitis. With its thin epithelial layers, the oral mucosa membrane might be sensitized to react to *Candida* and other virulence factors, thereby causing inflammation.

A chi-square test revealed that denture wearing was associated with the presence of atrophic glossitis (Table 4). The influence of salivation and denture wearing on *Candida* colony counts was evaluated using the Kruskal-

Wallis H test. The number of *C. albicans* colonies (Fig. 2) was found to be significantly higher in the hyposalivation group than in the normosalivation group. This result is in accordance with the findings of a previous study, which demonstrated an inverse association between the salivary flow rate and *C. albicans* counts in saliva [40–42]. However, no significant difference was observed in the amounts of *C. albicans* colonies between the denture wearers and denture non-wearers. *C. glabrata* exhibits superior cell surface hydrophobicity and a greater tendency to adhere to the surface of denture acrylic resin than other *Candida* species [43]. In addition, NAC shows a greater ability to form biofilms on the surface of dental acrylic resin in comparison to *C. albicans*, while the combination of *C. albicans* and *C. glabrata* shows the highest biofilm-forming ability [44]. The morphological differences and growth patterns exhibited by *C. albicans* and *C. glabrata* may allow these species to occupy the same location within the oral cavity with limited competition for space, thereby allowing their co-existence [33]. Thus, the presence of *Candida* on the denture surface is a prominent etiological factor for denture stomatitis [45]. However, there seems to be no correlation between the colonization of *Candida* on the mucosal surface of the denture base and that on the dorsum of the tongue. Although denture wearers with hyposalivation are considered at substantial risk for oral candidiasis, increased *C. albicans* colony counts on the dorsum of the tongue is essential for *Candida*-related atrophic glossitis.

Conclusion

C. albicans was associated with atrophic glossitis in xerostomia patients who had no systemic predisposing factors, indicating that *C. albicans* remains a treatment target for *Candida*-related atrophic glossitis.

Abbreviations

NAC: Non-*albicans Candida*; RSFR: Resting whole salivary flow rate; SSFR: Stimulated whole salivary flow rate

Acknowledgements

Not applicable.

Funding

No specific funding was received for this study.

Availability of data and materials

The datasets obtained or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

YN designed the study, performed clinical dental examinations of the patients and drafted the manuscript. SN, MR, KY, and IS performed clinical dental examinations of the patients and drafted the manuscript. NM performed the microbiological analyses. YY and HI participated in the analysis and interpretation of the data and drafted the manuscript. Statistical analyses were performed by KY and AT. All authors have read and approved the final version of this manuscript.

Ethics approval and consent to participate

All patients provided written informed consent. The Ethics Committee at Tsurumi University School of Dental Medicine approved this study (approval number 244, Aug. 19, 2004, and approval number 521, Mar. 21, 2008).

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department of Clinical Pathophysiology, Tsurumi University School of Dental Medicine, Tsurumi University Dental Hospital, 2-1-3 Tsurumi, Tsurumi-ku, Yokohama 230-8501, Japan. ²Department of Oral Microbiology, Tsurumi University, School of Dental Medicine, Yokohama, Japan. ³Kobayashi Dental Clinic, Niigata, Japan. ⁴Community Dentistry, Tsurumi University School of Dental Medicine, Yokohama, Japan. ⁵Department of Nutrition and Dietetics, Sagami Women's University, Sagami-hara, Japan. ⁶Department of Dentomaxillofacial Diagnosis and Treatment, Division of Oral and Maxillofacial Surgery, Graduate School of Kanagawa Dental University, Yokosuka, Japan. ⁷Department of Pathology, Tsurumi University School of Dental Medicine, Yokohama, Japan.

Received: 1 March 2017 Accepted: 7 December 2017

Published online: 16 December 2017

References

- Williams D, Lewis M. Pathogenesis and treatment of oral candidosis. *J Oral Microbiol.* 2011;3:5771. doi:10.3402/jom.v3i0.5771.
- Ellepola AN, Samaranyake LP. Oral candidal infections and antimycotics. *Crit Rev Oral Biol Med.* 2000;11(2):172–98.
- Reamy BV, Derby R, Bunt CW. Common tongue conditions in primary care. *Am Fam Physician.* 2010;81(5):627–34.
- Kurnatowska AJ. Search for correlation between symptoms and signs of changes in the oral mucosa and presence of fungi. *Mycoses.* 2001;44(9–10):379–82.
- Terai H, Shimahara M. Atrophic tongue associated with Candida. *J Oral Pathol Med.* 2005;34(7):397–400.
- Terai H, Shimahara M. Partial atrophic tongue other than median rhomboid glossitis. *Clin Exp Dermatol.* 2007;32(4):381–4.
- Kimori H, Yamamoto K, Yamachika S, Tsurumoto A, Kamikawa Y, Sasao M, Morito M, Saito I, Oshima T, Maeda N, et al. Factors associated with the presence of atrophic tongue in patients with dry mouth. *Gerodontology.* 2015;32(1):13–7.
- Shinozaki S, Moriyama M, Hayashida JN, Tanaka A, Maehara T, Ieda S, Nakamura S. Close association between oral Candida species and oral mucosal disorders in patients with xerostomia. *Oral Dis.* 2012;18(7):667–72.
- Hertel M, Schmidt-Westhausen AM, Strietzel FP. Local, systemic, demographic, and health-related factors influencing pathogenic yeast spectrum and antifungal drug administration frequency in oral candidiasis: a retrospective study. *Clin Oral Investig.* 2016;20(7):1477–86.
- Muadcheingka T, Tantivitayakul P. Distribution of Candida albicans and non-albicans Candida species in oral candidiasis patients: correlation between cell surface hydrophobicity and biofilm forming activities. *Arch Oral Biol.* 2015;60(6):894–901.
- Rodrigues CF, Silva S, Henriques M. Candida glabrata: a review of its features and resistance. *Eur J Clin Microbiol Infect Dis.* 2014;33(5):673–88.
- Samaranyake YH, Samaranyake LP. Candida krusei: biology, epidemiology, pathogenicity and clinical manifestations of an emerging pathogen. *J Med Microbiol.* 1994;41(5):295–310.
- Fidel PL Jr, Vazquez JA, Sobel JD. Candida glabrata: review of epidemiology, pathogenesis, and clinical disease with comparison to C. albicans. *Clin Microbiol Rev.* 1999;12(1):80–96.
- Coco BJ, Bagg J, Cross LJ, Jose A, Cross J, Ramage G. Mixed Candida albicans and Candida glabrata populations associated with the pathogenesis of denture stomatitis. *Oral Microbiol Immunol.* 2008;23(5):377–83.
- Vitali C, Bombardieri S, Jonsson R, Moutsopoulos HM, Alexander EL, Carsons SE, Daniels TE, Fox PC, Fox RI, Kassan SS, et al. Classification criteria for Sjögren's syndrome: a revised version of the European criteria proposed by the American-European Consensus Group. *Ann Rheum Dis.* 2002;61(6):554–8.
- Sreebny LM, Valdin A. Xerostomia. Part I: relationship to other oral symptoms and salivary gland hypofunction. *Oral Surg Oral Med Oral Pathol.* 1988;66(4):451–8.
- Okamoto MR, Kamoi M, Yamachika S, Tsurumoto A, Imamura T, Yamamoto K, Kadomatsu S, Saito I, Maeda N, Nakagawa Y. Efficacy of Fungiflora Y staining for the diagnosis of oral erythematous candidiasis. *Gerodontology.* 2013;30(3):220–5.
- Fujibayashi T, Sugai S, Miyasaka N, Hayashi Y, Tsubota K. Revised Japanese criteria for Sjögren's syndrome (1999): availability and validity. *Mod Rheumatol.* 2004;14:425–34.
- Williams DW, Lewis MA. Isolation and identification of Candida from the oral cavity. *Oral Dis.* 2000;6(1):3–11.
- Kimori H, Nakagawa Y, Yamamoto K, Oshima T. Establishing the cut-off point for the Candida swab test for daily oral care in dry mouth patients. *Oral Ther Pharmacol.* 2009;28(1):17–25.
- Tooyama H, Matsumoto T, Hayashi K, Kurashina K, Kurita H, Uchida M, Kasuga E, Honda T. Candida concentrations determined following concentrated oral rinse culture reflect clinical oral signs. *BMC Oral Health.* 2015;15:150.
- Neppelenbroek KH, Seo RS, Urban VM, Silva S, Dovigo LN, Jorge JH, Campanha NH. Identification of Candida species in the clinical laboratory: a review of conventional, commercial, and molecular techniques. *Oral Dis.* 2014;20(4):329–44.
- Powell HL, Sand CA, Rennie RP. Evaluation of CHROMagar Candida for presumptive identification of clinically important Candida species. *Diagn Microbiol Infect Dis.* 1998;32(3):201–4.
- Pfäller MA, Houston A, Coffmann S. Application of CHROMagar Candida for rapid screening of clinical specimens for Candida albicans, Candida tropicalis, Candida krusei, and Candida (Torulopsis) glabrata. *J Clin Microbiol.* 1996;34(1):58–61.
- Odds FC, Bernaerts R. CHROMagar Candida, a new differential isolation medium for presumptive identification of clinically important Candida species. *J Clin Microbiol.* 1994;32(8):1923–9.
- Beighton D, Ludford R, Clark DT, Brailsford SR, Pankhurst CL, Tinsley GF, Fiske J, Lewis D, Daly B, Khalifa N, et al. Use of CHROMagar Candida medium for isolation of yeasts from dental samples. *J Clin Microbiol.* 1995;33(11):3025–7.
- Campanha NH, Neppelenbroek KH, Spolidorio DM, Spolidorio LC, Pavarina AC. Phenotypic methods and commercial systems for the discrimination between C. albicans and C. dubliniensis. *Oral Dis.* 2005;11(6):392–8.
- Neppelenbroek KH, Campanha NH, Spolidorio DM, Spolidorio LC, Seo RS, Pavarina AC. Molecular fingerprinting methods for the discrimination between C. albicans and C. dubliniensis. *Oral Dis.* 2006;12(3):242–53.
- Sullivan DJ, Westerneng TJ, Haynes KA, Bennett DE, Coleman DC. Candida dubliniensis sp. nov.: phenotypic and molecular characterization of a novel species associated with oral candidosis in HIV-infected individuals. *Microbiology.* 1995;141(Pt 7):1507–21.
- Terai H, Shimahara M. Usefulness of culture test and direct examination for the diagnosis of oral atrophic candidiasis. *Int J Dermatol.* 2009;48(4):371–3.
- Aleva NA, Birman EG, Afonso W Jr, Chavasco JK, Paula CR, Ribeiro A, Pereira LJ. Erythematous candidosis in patients with complete dentures and HIV +/AIDS. *Mycoses.* 2007;50(5):407–11.
- Reichart PA, Samaranyake LP, Philipsen HP. Pathology and clinical correlates in oral candidiasis and its variants: a review. *Oral Dis.* 2000;6(2):85–91.
- Silva S, Henriques M, Hayes A, Oliveira R, Azeredo J, Williams DW. Candida glabrata and Candida albicans co-infection of an in vitro oral epithelium. *J Oral Pathol Med.* 2011;40(5):421–7.
- Sanita PV, Zago CE, Mima EG, Pavarina AC, Jorge JH, Machado AL, Vergani CE. In vitro evaluation of the enzymatic activity profile of non-albicans Candida species isolated from patients with oral candidiasis with or without diabetes. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2014;118(1):84–91.
- Tang SX, Moyes DL, Richardson JP, Blagojevic M, Naglik JR. Epithelial discrimination of commensal and pathogenic Candida albicans. *Oral Dis.* 2016;22(Suppl 1):114–9.

36. Villar CC, Zhao XR. *Candida albicans* induces early apoptosis followed by secondary necrosis in oral epithelial cells. *Mol Oral Microbiol*. 2010; 25(3):215–25.
37. Rhodus NL, Bloomquist C, Liljemark W, Bereuter J. A comparison of three methods for detecting *Candida albicans* in patients with Sjogren's syndrome. *Quintessence Int*. 1998;29(2):107–13.
38. Romagnoli P, Pimpinelli N, Mori M, Reichart PA, Eversole LR, Ficarra G. Immunocompetent cells in oral candidiasis of HIV-infected patients: an immunohistochemical and electron microscopical study. *Oral Dis*. 1997; 3(2):99–105.
39. Eversole LR, Reichart PA, Ficarra G, Schmidt-Westhausen A, Romagnoli P, Pimpinelli N. Oral keratinocyte immune responses in HIV-associated candidiasis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 1997; 84(4):372–80.
40. Torres SR, Peixoto CB, Caldas DM, Silva EB, Akiti T, Nucci M, de Uzeda M. Relationship between salivary flow rates and *Candida* counts in subjects with xerostomia. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2002; 93(2):149–54.
41. Epstein JB, Pearsall NN, Truelove EL. Quantitative relationships between *Candida albicans* in saliva and the clinical status of human subjects. *J Clin Microbiol*. 1980;12(3):475–6.
42. Navazesh M, Wood GJ, Brightman VJ. Relationship between salivary flow rates and *Candida albicans* counts. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 1995;80(3):284–8.
43. Luo G, Samaranayake LP. *Candida glabrata*, an emerging fungal pathogen, exhibits superior relative cell surface hydrophobicity and adhesion to denture acrylic surfaces compared with *Candida albicans*. *APMIS*. 2002; 110(9):601–10.
44. Pathak AK, Sharma S, Shrivastva P. Multi-species biofilm of *Candida albicans* and non-*Candida Albicans* *Candida* species on acrylic substrate. *J Appl Oral Sci*. 2012;20(1):70–5.
45. Altarawneh S, Bencharit S, Mendoza L, Curran A, Barrow D, Barros S, Preisser J, Loewy ZG, Gendreau L, Offenbacher S. Clinical and histological findings of denture stomatitis as related to intraoral colonization patterns of *Candida albicans*, salivary flow, and dry mouth. *J Prosthodont*. 2013;22(1):13–22.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

